

Distributed private randomness distillation

Dong Yang

China Jiliang University
University of Bergen

joint work [arXiv:1803.09989] with
Karol Horodecki (Univ. Gdansk) and Andreas Winter (Univ. Autònoma Barcelona)

Shanghai, 29 August 2018

1 Introduction

- Randomness
- BFW
- motivation

2 Results

- two-sided randomness distillation
- one-sided randomness distillation
- private randomness capacity

3 Summary

Randomness

- randomness has various applications
- classical world: pseudo randomness
- quantum mechanics: true randomness




Figure: $|+\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$, measurement basis $\{|0\rangle, |1\rangle\}$

private randomness





Figure: $|\Phi\rangle_{AE} = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$, measurement basis $\{|0\rangle, |1\rangle\}$

Local noise $\Phi_A = \frac{1}{2}(|0\rangle\langle 0| + |1\rangle\langle 1|)$ is useless by itself.


Berta/Fawzi/Wehner [IEEE Trans. Inf. Theory 60,
1168 (2014)]



$$\rho_{AE}^{\otimes n} \xrightarrow{U_{A \rightarrow KA'}} \sigma_{KA'E} \xrightarrow{\text{Tr}_{A'}} \sigma_{KE} \xrightarrow{M:|i\rangle_K} \sigma_{\hat{K}E} \stackrel{1-\epsilon}{\approx} \frac{1}{K} \sum_{i=1}^K |i\rangle\langle i| \otimes \rho_E^{\otimes n}$$


- question: $R_A = \sup \frac{\log K}{n}$, s.t. $n \rightarrow \infty$, $\epsilon \rightarrow 0$?
- answer: $R_A = \log |A| + [S(A|E)]_-$, where $[t]_- := \min\{0, t\}$

motivation



- a hidden point in BFW: a trusted but passive Bob
- idea: a trusted and **active** Bob?


our scenario



$$\psi_{ABE}^{\otimes n} \frac{w/o \text{ noise}}{w/o \text{ CC}} \rightsquigarrow \approx^{1-\epsilon} \frac{1}{K_A} \sum_{i=1}^{K_A} |i\rangle\langle i| \otimes \frac{1}{K_B} \sum_{j=1}^{K_B} |j\rangle\langle j| \otimes \psi_E^{\otimes n}$$

- a dual setting to distributed data compression (Slepian/Wolf) in classical IT whose natural dual setting **DOES NOT** exist classically, but **DOES** quantumly!
- the rate region of (R_A, R_B) , where $R_A = \frac{\log K_A}{n}$, $R_B = \frac{\log K_B}{n}$, s.t. $n \rightarrow \infty$, $\epsilon \rightarrow 0$?

warmup: entanglement swapping



Local noise can help!

idea




Figure: $S(A|B) > 0 > S(B|A)$. no communication, no noise.

Local noise can be created!

protocol for two-sided randomness distillation

no noise and no communication

$$\begin{aligned}R_A &\leq \log |A| - S(A|B)_+, \\R_B &\leq \log |B| - S(B|A)_+, \\R_A + R_B &\leq R_G = \log |AB| - S(AB),\end{aligned}$$

where $[t]_+ = \max\{0, t\}$.

free noise but no communication

$$\begin{aligned}R_A &\leq \log |A| - S(A|B), \\R_B &\leq \log |B| - S(B|A), \\R_A + R_B &\leq R_G = \log |AB| - S(AB).\end{aligned}$$

free noise and free communication

$$R_A \leq R_G,$$

$$R_B \leq R_G,$$

$$R_A + R_B \leq R_G = \log |AB| - S(AB),$$

free communication but no noise

$$R_A \leq \log |A| - \max\{S(B), S(AB)\},$$

$$R_B \leq \log |B| - \max\{S(A), S(AB)\},$$

$$R_A + R_B \leq R_G = \log |AB| - S(AB).$$




Figure: $S(A|B) > 0 > S(B|A)$. no comm. no noise, no comm. free noise, free comm. free noise, and free comm. no noise.

Remarks

- no cc, no/free noise is **dual to the Slepian-Wolf theorem!**
- local noise can boost randomness extraction
- **no bound** randomness state
- the rate regions are **tight** in
 - no cc, no noise
 - no cc, free noise
 - free cc, free noise.

tight for free cc, no noise ?

one-sided randomness distillation

- task: Bob helps Alice to extract randomness against Eve.
- the rate

$$R_A = \log |AB| - \inf_n \frac{1}{n} \max\{S(E'^{(n)}), S(B'^{(n)})\}$$

$$RCLOCC : \rho_{AB}^{\otimes n} \rightarrow \sigma_{A'^n B'^n}$$

- not strongly additive

$$R_A(\rho \otimes \sigma) > R_A(\rho) + R_A(\sigma)$$

- Bell state

$$R_A(\Phi) = \frac{3}{2}$$

- upper bound

$$\begin{aligned} R_A &\leq \log |AB| - \max\left\{\frac{1}{2}[E_r^\infty(\rho_{AB}) + S(AB)], S(AB)\right\} \\ &\leq \log |AB| - \frac{1}{2} \max\{S(A), S(B)\} \end{aligned}$$

private randomness capacity




Figure: server-client model: $|\phi\rangle_{AA'} \xrightarrow{\mathcal{N}: A' \rightarrow B} \rho_{AB}$

- task: Bob extract randomness secure against Eve.
- question: what is the maximal rate?
- in line with the standard model of transmitting information
- server-client structure in future quantum networks

private randomness capacity

Answer

- $\log d_B + \max_{|\phi\rangle_{AA'}} \{S(A) - S(\mathcal{N}_{A' \rightarrow B}(\phi_{AA'}))\}$!
- reverse coherent information of a channel
 $I_{rev}(\mathcal{N}) = \max_{|\phi\rangle_{AA'}} \{S(A) - S(\mathcal{N}_{A' \rightarrow B}(\phi_{AA'}))\}$

Remark

- single-letter formula, computable.
- $I_{rev}(\mathcal{N} \otimes \mathcal{M}) = I_{rev}(\mathcal{N}) + I_{rev}(\mathcal{M})$
[CMP266, 37 (2006)], [PRL102, 210501 (2009)]
- But its interpretation has been missing since then.

In contrast with coherent information

- $I_{coh}(\mathcal{N}) = \max_{|\phi\rangle_{AA'}} \{S(B) - S(\mathcal{N}_{A' \rightarrow B}(\phi_{AA'}))\}$
- $I_{coh}(\mathcal{M} \otimes \mathcal{N}) \neq I_{coh}(\mathcal{M}) + I_{coh}(\mathcal{N})$
- $Q(\mathcal{N}) = \sup \frac{1}{n} I_{coh}(\mathcal{N}^{\otimes n})$

Our scenario

distributed private randomness distillation

Results


- two-sided private randomness distillation
- one-sided private randomness distillation
- the private randomness capacity of a channel

Questions

- Q1. tight for free cc, no noise?
- Q2. is regularization in one-sided setting necessary?
- Q3. multipartite case, single-shot case, strong converse?

Thank you !

distributed data compression



- task: joint prob. distribution p_{XY} . Alice compresses her data at the rate R_X and Bob at R_Y . Charlie can recover the whole data reliably after receiving their compressed data.
- question: what is the compression rate region of (R_X, R_Y) ?
- answer: $R_X \geq H(X|Y)$, $R_Y \geq H(Y|X)$, $R_{XY} \geq H(XY)$
[Slepian-Wolf Theorem]

resource theory

allowed operations: restricted closed local operations and classical communication (RCLOCC)

- adding $|0\rangle$
- local unitary
- partial tracing
- local noise: $\frac{1}{2}(|0\rangle\langle 0| + |1\rangle\langle 1|)$
- classical communication: exchanging subsystems by a dephasing channel $\mathcal{N}(\rho) = \sum \langle i|\rho|i\rangle |i\rangle\langle i|$

In the standard picture, a state of one ideal randomness

$$\rho_{ABE} = \frac{1}{2}(|0\rangle\langle 0| + |1\rangle\langle 1|)_A \otimes \frac{1}{2}(|0\rangle\langle 0| + |1\rangle\langle 1|)_B \otimes \rho_E$$

in the dual picture, an ibit is

$$\alpha_{AA'BB'} = \frac{1}{4} \sum_{i,j,k,l=0}^1 |i\rangle\langle j|_A \otimes |k\rangle\langle l|_B \otimes U_{ik} \sigma_{A'B'} U_{jl}^\dagger$$

Quantum State Merging




Figure: $S(A|E) < 0$, classical communication $I(A : B)$, entanglement between A' and E' is $-S(A|E)$.

Local noise can be created from QST!